

ON GENERALIZED WHITEHEAD PRODUCTS

BRAYTON GRAY

Whitehead products have played an important role in unstable homotopy. They were originally introduced [Whi41] as a bilinear pairing of homotopy groups:

$$\pi_m(X) \otimes \pi_n(X) \rightarrow \pi_{m+n-1}(X) \quad m, n > 1.$$

This was generalized ([Ark62],[Coh57],[Hil59]) by constructing a map:

$$W: S(A \wedge B) \rightarrow SA \vee SB.$$

Precomposition with W defines a function on based homotopy classes:

$$[SA, X] \times [SB, X] \rightarrow [S(A \wedge B), X]$$

which is bilinear in case A and B are suspensions.

The case where A and B are Moore spaces was central to the work of Cohen, Moore and Neisendorfer ([CMN79]). In [Ani93] and in particular [AG95], this work was generalized. Much of this has since been simplified in [GT10], but further understanding will require a generalization from suspensions to co-H spaces.

The purpose of this work is to carry out and study such a generalization. Let $\mathcal{CO}$ be the category of simply connected co-H spaces and co-H maps. We define a functor:

$$\begin{aligned} \mathcal{CO} \times \mathcal{CO} &\rightarrow \mathcal{CO} \\ (G, H) &\rightarrow G \circ H \end{aligned}$$

and a natural transformation:

$$(1) \quad W: G \circ H \rightarrow G \vee H$$

generalizing the Whitehead product map. The existence of $G \circ H$ generalizes a result of Theriault [The03] who showed that the smash product of two simply connected co-associative co-H spaces is the suspension of a co-H space. We do not need the co-H spaces to be co-associative and require only one of them to be simply connected.* We call $G \circ H$ the Theriault product of G and H .

We summarize our results in the following theorems.

*In fact we can define $G \circ H$ for any two co-H spaces but require at least one of them to be either simply connected or a suspension in order to obtain the co-H space structure map on $G \circ H$.

Theorem 1. *There is a functor: $\mathcal{CO} \times \mathcal{CO} \rightarrow \mathcal{CO}$ given by*

$$(G, H) \rightarrow G \circ H$$

and equivalences in $\mathcal{CO}$

- (a) $(SX) \circ H \simeq X \wedge H$
- (b) $S(G \circ H) \simeq G \wedge H$
- (c) $(G_1 \vee G_2) \circ H \simeq G_1 \circ H \vee G_2 \circ H$

and homotopy equivalences:

- (d) $G \circ H \simeq H \circ G$
- (e) $(G \circ H) \circ K \simeq G \circ (H \circ K)$

Theorem 2. *There is a natural transformation:*

$$W: G \circ H \rightarrow G \vee H$$

which is the Whitehead product map (1) in case G and H are both suspensions. Furthermore, there is a homotopy equivalence:

$$G \times H \simeq G \vee H \cup_W C(G \circ H).$$

The next theorem concerns the inclusion of the fiber in certain standard fibration sequences [Gra71]:

$$\begin{aligned} G \times \Omega H &\xrightarrow{\iota_1} G \vee H \xrightarrow{\pi_2} H \\ \Omega G * \Omega H &\xrightarrow{\iota_2} G \vee H \rightarrow G \times H. \end{aligned}$$

Define $\text{ad}^n(H)(G)$ inductively by $\text{ad}^0(H)(G) = G$ and

$$\text{ad}^n(H)(G) = [\text{ad}^{n-1}(H)(G)] \circ H$$

and an iterated Whitehead product

$$\text{ad}^n: \text{ad}^n(H)(G) \rightarrow G \vee H$$

as the composition:

$$\text{ad}^n(H)(G) \xrightarrow{W} \text{ad}^{n-1}(H)(G) \vee H \xrightarrow{\text{ad}^{n-1} \vee 1} G \vee H \vee H \rightarrow G \vee H.$$

Theorem 3. *Suppose G and H are simply connected co- H spaces. Then there are homotopy equivalences:*

- (a) $G \times \Omega H \simeq \bigvee_{n \geq 0} \text{ad}^n(H)(G)$ where ι_1 corresponds to ad^n on the appropriate factor
- (b) $\Omega G * \Omega H \simeq \bigvee_{\substack{i \geq 0 \\ j \geq 1}} \text{ad}^i(H)(\text{ad}^j(G)(G))$ where ι_2 corresponds to $\text{ad}^i(\text{ad}^j)$ on to the appropriate factor
- (c) $S\Omega G \simeq \bigvee_{n \geq 0} \text{ad}^n(G)(G)$ where the composition $S\Omega G \rightarrow S\Omega G \vee S\Omega G \rightarrow G \vee G$ corresponds to the appropriate iterated Whitehead product on each factor.

It should be pointed out that the equivalence (c) generalizes the result of Theriault [The03, 1.1] where it is shown that a simply connected co-associative co-H space decomposes

$$\Sigma\Omega G \simeq \bigvee_{n \geq 1} M_n$$

for some spaces M_n , which are not further decomposed.

Theorem 4. *Suppose X is finite dimensional and $f: SX \rightarrow G \vee H$ then f is the sum of the projections onto G and H and a finite sum of iterated Whitehead products.*

Throughout this work we will assume that all spaces are of the homotopy type of a CW complex. All homology and cohomology will be with a field of coefficients. We will often show that a map between simply connected CW complexes is a homotopy equivalence by showing that it induces an isomorphism in homology with an arbitrary field of coefficients, without further comment.

Section 1 will be devoted to some general remarks about telescopes and we will construct the Theriault product in section 2. Theorem 1 will follow from 2.3, 2.5 and 2.7. The functor in Theorem 2 is defined after 3.2 and the equivalence follows from 3.8. The proof of the first part of Theorem 3 occurs just prior to 3.5 and the rest follows from 3.5 Theorem 4 follows from 3.7.

1.

In this section we will discuss some general properties of telescopes of a self map $e: G \rightarrow G$ where G is a co-H space. We do not assume that e is idempotent. We will call e a quasi-idempotent if the induced homomorphism in homology satisfies the equation:

$$(e_*)^2 = -e_*$$

where u is a unit. We construct two telescopes:

$$\begin{aligned} T(e): G &\xrightarrow{e} G \xrightarrow{e} G \rightarrow \dots \\ T(1+e): G &\xrightarrow{1+e} G \xrightarrow{1+e} G \rightarrow \dots \end{aligned}$$

and a map:

$$\Gamma: G \rightarrow G \vee G \xrightarrow{\Gamma_1 \vee \Gamma_2} T(e) \vee T(1+e).$$

Proposition 1.1. *If G is simply connected and e is a quasi-idempotent, Γ is a homotopy equivalence. Furthermore $H_*(T(e)) = \text{im } e_*$ and $\tilde{H}_*(T(1+e)) \cong \ker e_*$.*

Proof. Suppose $\Gamma_*(\xi) = 0$. Since $(\Gamma_1)_*(\xi) = 0$ $(e_*)^k(\xi) = 0$ for some k , so $e_*(\xi) = 0$. Since $(\Gamma_2)_*(\xi) = 0$, $(1+e)_*^k(\xi) = 0$. But $(1+e)_*^2 = (1+e)_*$, so $\xi = -e_*(\xi) = 0$. Clearly Γ_* is onto. Moreover, $H_*(T(e)) \cong \text{im } e_*$ and $\tilde{H}_*(T(1+e)) \cong \text{im}(1+e_*) = \ker e_*$.

Corollary 1.2. *Suppose G is a simply connected co- H space and $A \subset G$ is a retract of G . Let e be the composition:*

$$G \xrightarrow{r} A \xrightarrow{i} G.$$

Then $T(e) \simeq A$, $T(1 - e) \simeq G/A$, and the identity map of $T(e)$ can be factored:

$$T(e) \xrightarrow{\xi} A \xrightarrow{i} G \xrightarrow{r} A \xrightarrow{\eta} T(e)$$

where ξ and η are inverse homotopy equivalences.

Proof. The telescope $T(e)$ and A are both simply connected and there are maps $T(e) \rightarrow A$ and $A \rightarrow T(e)$ making A a retract of $T(e)$, and these maps are homotopy equivalences. By the Van Kampen theorem $G/A \simeq G \cup CA$ is simply connected. Since $1 - e: G \rightarrow G$ factors through the projection $\pi: G \rightarrow G/A$, we can factor the identity map up to homotopy

$$G \rightarrow G \vee G \xrightarrow{r \vee \pi} A \vee G/A \rightarrow G$$

and hence $G \simeq A \vee G/A$. The factorization of the identity map of $T(e)$ is obtained by replacing each space by a telescope where the three in the center are constant.

Now consider two maps $f_1, f_2: X \rightarrow X$.

Proposition 1.3. $T(f_1 f_2) \simeq T(f_2 f_1)$.

Proof. We define maps between the telescopes:

$$\begin{array}{ccccccc} X & \xrightarrow{f_1 f_2} & X & \xrightarrow{f_1 f_2} & X & \longrightarrow & \cdots \\ f_2 \downarrow & & f_2 \downarrow & & f_2 \downarrow & & \\ X & \xrightarrow{f_2 f_1} & X & \xrightarrow{f_2 f_1} & X & \longrightarrow & \cdots \\ f_1 \downarrow & & f_1 \downarrow & & f_1 \downarrow & & \\ X & \xrightarrow{f_1 f_2} & X & \xrightarrow{f_1 f_2} & X & \longrightarrow & \cdots \end{array}$$

The composition is the shift map which is a homotopy equivalence.

2.

In this section we will consider a pair of co- H spaces in which at least one is simply connected. Let G and H be two such co- H spaces with their structure determined by maps $\nu_1: G \rightarrow S\Omega G$ and $\nu_2: H \rightarrow S\Omega H$ each of which is a right inverse to the respective evaluation maps (See [Gan70]), which we label as ϵ_1, ϵ_2 . We define self maps of $S(\Omega G \wedge \Omega H)$ as follows:

$$e_1: S(\Omega G \wedge \Omega H) \xrightarrow{\epsilon_1 \wedge 1} G \wedge \Omega H \xrightarrow{\nu_1 \wedge 1} S(\Omega G \wedge \Omega H)$$

$$e_2: S(\Omega G \wedge \Omega H) \xrightarrow{1 \wedge \epsilon_2} \Omega G \wedge H \xrightarrow{1 \wedge \nu_2} S(\Omega G \wedge \Omega H);$$

here we freely move the suspension coordinate to wherever it is needed. Clearly e_1 and e_2 are idempotents but $e_1 e_2$ is not an idempotent; however

it is a quasi-idempotent. In fact by swapping coordinates, one can see that $S(e_1e_2)$ is homotopic to the negative of the composition:

$$\bar{e}: S^2(\Omega G \wedge \Omega H) \xrightarrow{\epsilon_1 \wedge \epsilon_2} G \wedge H \xrightarrow{\nu_1 \wedge \nu_2} S^2(\Omega G \wedge \Omega H).$$

Since $\bar{e}$ is clearly an idempotent $S(e_1e_2) \circ S(e_1e_2) \sim \bar{e}^2 = \bar{e} \sim -S(e_1e_2)$, so $(e_1e_2)_*^2 = -(e_1e_2)_*$.

Now assuming that one of G, H is simply connected, it follows that $\Omega G \wedge \Omega H$ is connected, so $S(\Omega G \wedge \Omega H)$ is simply connected. Consequently

Proposition 2.1. *If one of G and H is simply connected, there is a homotopy equivalence:*

$$S(\Omega G \wedge \Omega H) \simeq T(e_1e_2) \vee T(1 + e_1e_2).$$

Let

$$\theta: S(\Omega G \wedge \Omega H) \rightarrow T(e_1e_2)$$

be the projection and

$$\psi: T(e_1e_2) \rightarrow S(\Omega G \wedge \Omega H)$$

be the unique right inverse to θ which projects trivially onto $T(1 + e_1e_2)$. These maps determine a co-H space structure on $T(e_1e_2)$.

Definition 2.2. $G \circ H = T(e_1e_2)$.

Proposition 2.3. *Given co-H maps $f: G \rightarrow G'$ and $g: H \rightarrow H'$, there is an induced co-H map*

$$f \circ g: G \circ H \rightarrow G' \circ H'$$

making $G \circ H$ a functor of two variables. In addition there are equivalences of co-H spaces.

- (a) $SX \circ H \simeq X \wedge H$
- (b) $S(G \circ H) \simeq G \wedge H$
- (c) $S^1 \circ H \simeq H$

and there is a homotopy equivalence $G \circ H \simeq H \circ G$.

Proof. Since f and g are co-H maps, the squares

$$\begin{array}{ccc} G & \xrightarrow{\nu_1} & S\Omega G \\ f \downarrow & S\Omega f \downarrow & \\ G' & \xrightarrow{\nu'_1} & S\Omega G' \end{array} \quad \begin{array}{ccc} H & \xrightarrow{\nu_2} & S\Omega H \\ g \downarrow & S\Omega g \downarrow & \\ H' & \xrightarrow{\nu'_2} & S\Omega H' \end{array}$$

commute up to homotopy. It follows that f and g induce maps that commute with e_1 and e_2 and hence with the equivalences of 2.1, θ and ψ . For part a, observe that the composition e_2e_1 factors:

$$\begin{array}{ccccccc} S(\Omega SX \wedge \Omega H) & \xrightarrow{\epsilon_1 \wedge 1} & SX \wedge \Omega H & \xrightarrow{S\epsilon \wedge 1} & S\Omega SX \wedge \Omega H & \xrightarrow{1 \wedge \epsilon_2} & \Omega SX \wedge H \xrightarrow{1 \wedge \nu_2} S(\Omega SX \wedge \Omega H) \\ & & & \searrow 1 \wedge G & & \nearrow L \wedge 1 & \\ & & & & X \wedge H & & \end{array}$$

where $(1 \wedge \epsilon)(\epsilon_1 \wedge 1)$ is a right universe to $(1 \wedge \nu_2)(\iota \wedge 1)$. Thus we can apply 1.2 to see that $SX \circ H \simeq X \wedge H$ with co-H structure given by the composite $(1 \wedge \nu_2)(\iota \wedge 1)$. This is precisely the co-H structure induced by ν_2 . Part b follows since $S(G \circ H)$ is the telescope of $\bar{e}$ with co-H structure given by $\nu_1 \wedge \nu_2$. Part c is a special case of part a: The last statement follows directly from 1.3.

For the associativity assertion in Theorem 1, it will be convenient to describe an alternative definition of $G \circ H$. For this we assume that G is a retract of a space SX and H a retract of SY :

$$G \xrightarrow{\nu_1} SX \xrightarrow{\epsilon_1} G \quad H \xrightarrow{\nu_2} SY \xrightarrow{\epsilon_2} H.$$

We can then replace the telescope in the definition by the telescope of the composition:

$$T: SX \wedge Y \xrightarrow{1 \wedge \epsilon_2} X \wedge H \xrightarrow{1 \wedge \nu_2} SX \wedge Y \xrightarrow{\epsilon_1 \wedge 1} G \wedge Y \xrightarrow{\nu_1 \wedge 1} SX \wedge Y.$$

The co-H structures defined by these maps are equivalent to the structures defined by

$$\tilde{\nu}_2: G \xrightarrow{\nu_1} SX \xrightarrow{S\tilde{\epsilon}_1} S\Omega X \quad \tilde{\nu}_1: H \xrightarrow{\nu_2} SY \xrightarrow{S\Omega\tilde{\epsilon}_2} S\Omega H$$

and we have a homotopy commutative ladder:

$$\begin{array}{ccccccccc} SX \wedge Y & \xrightarrow{1 \wedge \epsilon_2} & X \wedge H & \xrightarrow{1 \wedge \nu_2} & SX \wedge Y & \xrightarrow{\epsilon_1 \wedge 1} & G \wedge Y & \xrightarrow{\nu_2 \wedge 1} & SX \wedge Y \\ \downarrow S\tilde{\epsilon}_1 \wedge \tilde{\epsilon}_2 & & \downarrow \hat{\epsilon}_1 \wedge 1 & & \downarrow S\tilde{\epsilon}_1 \wedge \tilde{\epsilon}_2 & & \downarrow 1 \wedge \tilde{\epsilon}_2 & & \downarrow S\tilde{\epsilon}_1 \wedge \tilde{\epsilon}_2 \\ S\Omega G \wedge \Omega H & \xrightarrow{1 \wedge \epsilon} & \Omega G \wedge H & \xrightarrow{1 \wedge \tilde{\nu}_2} & S\Omega G \wedge \Omega H & \xrightarrow{\epsilon \wedge 1} & G \wedge \Omega H & \xrightarrow{\tilde{\nu}_1 \wedge 1} & S\Omega G \wedge \Omega H. \end{array}$$

Hence we have a commutative diagram

$$\begin{array}{ccc} SX \wedge Y & \xrightarrow{\quad} & T \vee \bar{T} \\ S\tilde{\epsilon}_1 \wedge \tilde{\epsilon}_2 \downarrow & & \alpha \vee \beta \downarrow \\ S\Omega G \wedge \Omega H & \longrightarrow & G \circ H \vee \text{Tel}(1 + e_1 e_2) \end{array}$$

where $\bar{T}$ is the telescope defined by $1 + (\nu_1 \wedge 1)(\epsilon_1 \wedge 1)(1 \wedge \nu_2)(1 \wedge \epsilon_2)$. The map $\alpha: T \rightarrow G \circ H$ is compatible with the homotopy equivalence $ST \simeq G \wedge H \simeq S(G \circ H)$ so is itself a homotopy equivalence. Now choose a right homotopy inverse for the map $SX \wedge Y \rightarrow T$ which projects trivially to $\bar{T}$. It's composite with $S\tilde{\epsilon}_1 \wedge \tilde{\epsilon}_2$ will then project trivially to $T(1 + e_1 e_2)$. Hence we have a homotopy commutative diagram:

$$\begin{array}{ccccc} T & \xrightarrow{\quad} & SX \wedge Y & \xrightarrow{\quad} & T \\ \alpha \downarrow & & S\tilde{\epsilon}_1 \wedge \tilde{\epsilon}_2 \downarrow & & \alpha \downarrow \\ G \circ H & \longrightarrow & S\Omega G \wedge \Omega H & \longrightarrow & G \circ H \end{array}$$

and consequently the co-H structure on T is compatible under α with the co-H structure on $G \circ H$. We have proven

Proposition 2.4. *Suppose G is represented as a retract of SX and H a retract of SY . Then $G \circ H$ is homotopy equivalent to the telescope T of the composition.*

$$SX \wedge Y \rightarrow G \wedge Y \rightarrow SX \wedge Y \rightarrow X \wedge H \rightarrow SX \wedge Y$$

as co- H spaces, where the co- H structure on T is given by the equivalence $SX \wedge Y \simeq T \vee \overline{T}$.

Proposition 2.5. $(G_1 \vee G_2) \circ H \simeq G_1 \circ H \vee G_2 \circ H$ as co- H spaces.

Proof. Write G_i as a retract of SX_i , $i = 1, 2$. Then $G_1 \vee G_2$ is a retract of $S(X_1 \vee X_2)$. Thus the telescope for $(G_1 \vee G_2) \circ H$ is at each point the wedge of the telescopes for $G_1 \circ H$ and $G_2 \circ H$.

At this point we will apply 2.4 to prove theorem 1 part e, the associativity formula. We will make repeated use of

Lemma 2.6. *There is a homotopy commutative square*

$$\begin{array}{ccc} S(G \circ H) & \xrightarrow{S\psi} & S^2X \wedge Y \\ \simeq \downarrow & \nearrow \nu_1 \wedge \nu_2 & \downarrow S\theta \\ G \wedge H & \xrightarrow{\simeq} & S(G \circ H). \end{array}$$

Proof. $G \wedge H$ is a retract of $S^2X \wedge Y$, so we may apply 1.2.

Proposition 2.7. $(G \circ H) \circ K \simeq G \circ (H \circ K)$.

Proof. We suppose that G , H , and K are presented by retractions

$$\begin{array}{ccc} G & \xrightarrow{\psi_1} & SX \xrightarrow{\theta_1} G \\ H & \xrightarrow{\psi_2} & SY \xrightarrow{\theta_2} H \\ K & \xrightarrow{\psi_3} & SZ \xrightarrow{\theta_3} K \end{array}$$

and we then construct retractions for $G \circ H$ and $H \circ K$:

$$\begin{array}{ccc} G \circ H & \xrightarrow{\psi_3} & SX \wedge Y \xrightarrow{\theta_3} G \circ H \\ H \circ K & \xrightarrow{\psi_4} & SY \wedge Z \xrightarrow{\theta_4} H \circ K. \end{array}$$

Using these we construct retractions for $G \circ (H \circ K)$ and $(G \circ H) \circ K$

$$\begin{array}{ccc} (G \circ H) \circ K & \xrightarrow{\psi} & S(X \wedge Y \wedge Z) \xrightarrow{\theta} (G \circ H) \circ K \\ G \circ (H \circ K) & \xrightarrow{\psi'} & S(X \wedge Y \wedge Z) \xrightarrow{\theta'} (G \circ (H \circ K)). \end{array}$$

We will show that $\theta'\psi: (G \circ H) \circ K \rightarrow G \circ (H \circ K)$ is a homotopy equivalence. By 2.6, we have homotopy commutative diagrams:

$$\begin{array}{ccccc}
 & (G \circ H) \wedge K & & G \wedge (H \circ K) & \\
 & \nearrow \simeq & \searrow \psi_4 \wedge \psi_3 & \swarrow \psi_1 \wedge \psi_5 & \searrow \simeq \\
 S((G \circ K) \circ K) & \xrightarrow{S\psi} & S^2 X \wedge Y \wedge Z & \xrightarrow{S\theta'} & S(G \circ (H \circ K)).
 \end{array}$$

Suspending and applying 2.6 again we obtain a homotopy commutative diagram:

$$\begin{array}{ccccc}
 & G \wedge H \wedge K & & & \\
 & \swarrow \simeq & \downarrow \psi_1 \wedge \psi_2 \wedge \psi_3 & \searrow \simeq & \\
 S(G \circ H) \wedge K & & & & SG \wedge (H \circ K) \\
 \searrow S(\psi_4 \wedge \psi_3) & & \downarrow & \swarrow S(\psi_1 \wedge \psi_5) & \\
 & S^3(X \wedge Y \wedge Z) & & &
 \end{array}$$

from these diagrams it follows that $S^2(\theta'\psi)$ is a homotopy equivalence and hence $\theta'\psi$ is as well.

3.

In this section we generalize the clutching construction [Gra88, Proposition 1] for fibrations over a suspension to fibrations over a co-H space. This allows for the decomposition results in theorems 2 and 3.

Proposition 3.1. *Suppose $F \rightarrow E \rightarrow G$ is a fibration where G is a co-H space. Then $E/F \simeq G \rtimes F$.*

Proof. In the case $G = SX$, we have by [Gra88, Proposition 1]

$$E \simeq F \cup_{\theta} (CX) \times F.$$

So $E/F \simeq SX \rtimes F$. It is easy to construct a map $G \rtimes F \rightarrow E/F$ in general. Consider the sequence of pull backs:

$$\begin{array}{ccccc}
 F & \xlongequal{\quad} & F & \xlongequal{\quad} & F \\
 \downarrow & & \downarrow & & \downarrow \\
 E & \longrightarrow & E' & \longrightarrow & E \\
 \downarrow & & \downarrow & & \downarrow \\
 G & \xrightarrow{\nu} & S\Omega G & \xrightarrow{\epsilon} & G.
 \end{array}$$

Then we consider the composite:

$$E/F \rightarrow E'/F \simeq S\Omega G \rtimes F \rightarrow G \rtimes F$$

where the middle equivalence follows since the base is a suspension. Showing that the composite is a homotopy equivalence will take some work.

Since $\nu\epsilon: S\Omega G \rightarrow S\Omega G$ is an idempotent, we can decompose $S\Omega G$:

$$S\Omega G \simeq G \vee G'.$$

We now observe that we can construct a quasifibration model for a fibration over a one point union. Suppose we have such a fibration

$$\begin{array}{ccccc} E_A & \longrightarrow & \tilde{E} & \longleftarrow & E_B \\ \downarrow & & \downarrow & & \downarrow \\ A & \longrightarrow & A \vee B & \longleftarrow & B \end{array}$$

with pull backs E_A and E_B and fiber F . Then we can construct

$$E_A \cup_F E_B$$

the union of E_A and E_B with the subspace F identified. Then

$$\begin{array}{ccc} E_A \cup_F E_B & \xrightarrow{\phi} & \tilde{E} \\ \downarrow & & \downarrow \\ A \vee B & \xlongequal{\quad} & A \vee B \end{array}$$

ϕ is a homotopy equivalence. In our case $S\Omega G \simeq G \vee G'$ and $E_G = E$, $E_{G'} = G' \times F$, so

$$\begin{array}{c} E' \simeq E \cup_F G' \times F \\ \downarrow \\ G \vee G' \end{array}$$

is a quasifiber by [DT58, 2.10]. On the other hand $E'/F \simeq S\Omega G \rtimes F \simeq G \rtimes F \vee G' \rtimes F$ while $E \cup_F G' \rtimes F \simeq E/F \vee G' \rtimes F$. Since the map between E' and $E \cup_F G' \rtimes F$ is a map over $G \vee G'$ we see that $E/F \simeq G \rtimes F$.

Corollary 3.2. $S\Omega G \simeq G \rtimes \Omega G$.

Proof. Apply 3.1 to the path space fibration over G .

Proof of theorem 2. Construction: We now describe our generalization of the Whitehead product. Suppose G and H are co-H spaces and one of them is simply connected. The Whitehead product:

$$W: G \circ H \rightarrow G \vee H.$$

is then defined as the composition:

$$G \circ H \xrightarrow{\psi} S\Omega G \wedge \Omega H \simeq \Omega G * \Omega H \xrightarrow{\omega} G \vee H$$

where ω is the inclusion of the fiber in the fibration sequence:

$$\Omega G * \Omega H \xrightarrow{\omega} G \vee H \rightarrow G \times H.$$

Clearly ψ and ω are natural transformations, so W is as well.

Before we prove the homotopy equivalence in theorem 2, we need to establish some results in theorem 3. We begin by constructing maps:

$$ad^n: ad^n(H)(G) \rightarrow G \vee H$$

inductively. For $n = 0$ this is just the inclusion of G in $G \vee H$. For $n > 0$ we define ad^n as the composition:

$$\begin{aligned} ad^n(H)(G) &= (ad^{n-1}(H)(G)) \circ H \\ &\rightarrow (ad^{n-1}(H)(G)) \vee H \rightarrow (G \vee H) \vee H = G \vee H. \end{aligned}$$

Next we calculate the effect of ad^n in loop space homology:

$$\Omega(ad^n)_*: H_*(\Omega(ad^n(H)(G))) \rightarrow H_*(\Omega(G \vee H))$$

To do this we need some notation. For each co-H space G , write $\sigma^{-1}: \tilde{H}_r(G) \rightarrow \tilde{H}_{r-1}(\Omega G)$ for the composition:

$$\tilde{H}_r(G) \xrightarrow{\nu_+} \tilde{H}_r(S\Omega G) \simeq \tilde{H}_{r-1}(\Omega G).$$

Let $\{x_i\}$ be a basis for $\tilde{H}_*(G)$. Then $H_*(\Omega G)$ is the tensor algebra on the classes $\{\sigma^{-1}(x_i)\}$. Given two classes $x \in \tilde{H}_r(G)$, $y \in \tilde{H}_s(H)$ we will write

$$x \circ y \in \tilde{H}_{r+s-1}(G \circ H)$$

for the class that corresponds to $x \triangle y \in \tilde{H}_{r+s}(G \wedge H)$ under the isomorphism:

$$\tilde{H}_{r-1}(G \circ H) \cong \tilde{H}_r(S(G \circ H)) \simeq \tilde{H}_r(G \wedge H).$$

Then the classes $\{x_i \circ y_j\}$ form a basis for $\tilde{H}_*(G \circ H)$ where $\{x_i\}$ and $\{y_i\}$ respectively are bases for $\tilde{H}_*(G)$ and $\tilde{H}_*(H)$.

Proposition 3.3. $(\Omega W)_*(\sigma^{-1}(x \circ y)) = \pm [\sigma^{-1}x, \sigma^{-1}y]$ where

$$(\Omega W)_*: H_*(\Omega(G \circ H)) \rightarrow H_*(\Omega(G \vee H)).$$

Proof. By lemma 2.6

$$\psi_*(x \circ y) = e \triangle \sigma^{-1}(x) \triangle \sigma^{-1}(y) \in H_*(S\Omega G \wedge \Omega H)$$

so

$$(\Omega\psi)_*(\sigma^{-1}(x \circ y)) = \sigma^{-1}(x) \triangle \sigma^{-1}(y) \in H_*(\Omega G \wedge \Omega H)$$

regarded as a submodule of $H_*(S\Omega G \wedge \Omega H)$. It now suffices to evaluate the composition:

$$\Omega G \wedge \Omega H \rightarrow \Omega(S\Omega G \wedge \Omega H) \xrightarrow{\Omega\xi} \Omega(\Omega G * \Omega H) \rightarrow \Omega(G \vee H)$$

where ξ is the standard homotopy equivalence $SX \wedge Y \simeq X * Y$.

Lemma 3.4. *The composition:*

$$\Omega G \wedge \Omega H \rightarrow \Omega S(\Omega G \wedge \Omega H) \xrightarrow{\Omega\xi} \Omega(\Omega G * \Omega H) \rightarrow \Omega(G \vee H)$$

carries $\sigma^{-1}(x) \triangle \sigma^{-1}(y) \in H_*(\Omega G \wedge \Omega H)$ to $\pm [\sigma^{-1}(x), \sigma^{-1}(y)]$.

Proof. We first need to describe the homotopy equivalence

$$SX \wedge Y \xrightarrow{\xi} X * Y.$$

Here we write points of the join as $tx + (1-t)y$, $0 \leq t \leq 1$. So $X * Y$ is the quotient of $X \times I \times Y$ given by the identifications $(x, 0, y) \sim (x', 0, y)$ and $(x, 1, y) \sim (x, 1, y')$. Then ξ is given by the formula:

$$\xi(t, x, y) = \begin{cases} (*, 1-3t, y) & 0 \leq 3t \leq 1 \\ (x, 3t-1, y) & 1 \leq 3t \leq 2 \\ (x, 3-3t, *) & 2 \leq 3t \leq 3 \end{cases}.$$

The map $\Omega X * \Omega Y \xrightarrow{\omega} X \vee Y$ is given by

$$(\omega_1, t, \omega_2) \rightarrow \begin{cases} \omega_1(2t) & 0 \leq 2t \leq 1 \\ \omega_2(2-2t) & 1 \leq 2t \leq 2 \end{cases}.$$

Combining these we get

$$S\Omega X * \Omega Y \xrightarrow{\xi} \Omega X * \Omega Y \xrightarrow{\omega} X \vee Y$$

with a 6-part formula:

$$(t, \omega_1, \omega_2) = \begin{cases} (*, \omega_2(6t)) & 0 \leq 6t \leq 1 \\ * & 1 \leq 6t \leq 2 \\ (\omega_1(6t-2), *) & 2 \leq 6t \leq 3 \\ (*, \omega_2(4-6t)) & 3 \leq 6t \leq 4 \\ * & 4 \leq 6t \leq 5 \\ (\omega_1(6-6t), *) & 5 \leq 6t \leq 6 \end{cases}$$

so the adjoint takes the pair (ω_1, ω_2) to the product of loops $\omega_1^{-1}\omega_2^{-1}\omega_1\omega_2$. The effect of this on a primitive element is the graded commutator.

Now the iterated circle product $ad^n(H)(G)$ has homology generated by classes of the form

$$(\dots((x \circ y_1) \circ y_2) \cdots \circ y_n)$$

where $x \in \tilde{H}_*(G)$ and $y_i \in \hat{H}_*(H)$. By 3.3

$$(\Omega W)_* (\sigma^{-1}(\dots(x \circ y_1) \circ y_2) \cdots \circ y_n)$$

is $\pm$ the graded commutator

$$[\dots [[\sigma^{-1}(x), \sigma^{-1}(y_1)], \sigma^{-1}(y_2)] \dots \sigma^{-1}(y_n)]$$

where the classes x and y_i are thought of as classes in $\tilde{H}_*(G \vee H)$.

Proof of theorem 3 part (a): Now let $G_* = \tilde{H}_*(G)$ and $H_* = \tilde{H}_*(H)$. Let $L(G_* \oplus H_*)$ be the free Lie algebra generated by G_* and H_* , and $L(H_*)$ the free Lie algebra generated by H_* . Then Neisendorfer has analyzed the kernel

$$L(G_* \vee H_*) \rightarrow L(H_*)$$

([Nei09, 8.7.4]). He has shown that this is the free Lie algebra

$$L \left(\bigoplus_{n \geq 0} \text{ad}^n(H_*)(G_*) \right)$$

The universal enveloping algebra is thus the tensor algebra generated by the elements $\text{ad}^n(H_*)(G_*)$ for $n \geq 0$. Consequently the fiber of the projection $\Omega(G \vee H) \rightarrow \Omega H$ is

$$\Omega \left(\bigvee_{n \geq 0} \text{ad}^n(H)(G) \right)$$

and this is homotopy equivalent to $\Omega(G \rtimes \Omega H)$ and the map

$$\bigvee_{n \geq 0} \text{ad}^n(H)(G) \rightarrow G \vee H$$

which factors through $G \rtimes \Omega H$ establishes the homotopy equivalence in theorem 3.

Corollary 3.5. (a) *If G is simply connected, $S\Omega G \simeq \bigvee_{n \geq 0} \text{ad}^n(G)(G)$*
 (b) *if both G and H are simply connected*

$$\Omega G * \Omega H \simeq \bigvee_{\substack{i \geq 0 \\ j \geq 1}} \text{ad}^j(H) (\text{ad}^i(G)(G)) .$$

Proof. For part (a) apply 3.2 and theorem 3 part a. For part (b), we expand

$$\begin{aligned} \Omega G * \Omega H &\simeq (S\Omega G) \wedge \Omega H \\ &\simeq \left[\bigvee_{i \geq 0} (\text{ad}^i(G)(G)) \right] \wedge \Omega H \\ &\simeq \bigvee_{i \geq 0} [\text{ad}^i(G)(G) \wedge \Omega H] \\ &\simeq \bigvee_{\substack{i \geq 0 \\ j \geq 1}} \text{ad}^j(H) (\text{ad}^i(G)(G)) \end{aligned}$$

using 2.5 and theorem 3 part a.

Given a Theriault product $P = G_1 \circ \cdots \circ G_s$ with some fixed association, let us write $\ell(P) = s$ for the length of P .

Theorem 3.6. *Suppose G and H are both simply connected co- H spaces and $k \geq 1$. Then there is a locally finite collection of iterated Theriault products $\{P_\alpha(k)\}$ of length ℓ_α and iterated Whitehead product maps:*

$$\omega_\alpha(k): P_\alpha(k) \rightarrow G \vee H$$

such that

$$\Omega(G \vee H) \simeq \Omega \left(\bigvee_{\ell_\alpha > k} P_\alpha(k) \right) \times \Omega \left(\prod_{\ell_\alpha \leq k} P_\alpha(k) \right)$$

and the factors of the righthand side are mapped to the lefthand side by the $\omega_\alpha(k)$.

Proof. For $k = 1$ we use the decomposition:

$$\Omega(G \vee H) \simeq \Omega(\Omega G * \Omega H) \times \Omega(G \times H)$$

where $\Omega G * \Omega H$ is a boquet of iterated Theriault products of length at least 2 by 3.5(b). Now we proceed by induction on k . Among the finite list of products $P_\alpha(k)$ of length $k + 1$, choose one which we label P . Then

$$\begin{aligned} \Omega \left(\bigvee_{\ell_\alpha > k} P_\alpha \right) &\simeq \Omega \left(P \vee \bigvee_{\substack{\ell_\alpha > k \\ P_\alpha \neq P}} P_\alpha \right) \\ &\simeq \Omega P \times \Omega \left(\bigvee_{\substack{\ell_\alpha > k \\ P_\alpha \neq P}} P_\alpha \rtimes \Omega P \right). \end{aligned}$$

The second factor has one less product of length $k + 1$. If we repeat this process once for each $P_\alpha(k)$ of length $k + 1$, we obtain:

$$\Omega \left(\bigvee_{\ell(P_\alpha) > k} P_\alpha \right) = \Omega P_j \times \cdots \times \Omega P_m \times \Omega \left(\bigvee_{\ell(P_\alpha) > k+1} P_\alpha(k+1) \right).$$

Now add the $P_1 \dots P_m$ to the list of P_α with $\ell(P_\alpha) \leq k$ to obtain all $P_\alpha(k+1)$ with length $\leq k + 1$.

Corollary 3.7. *Suppose X is a finite dimensional co-associative co-H space and $f: X \rightarrow G \vee H$ where G and H are simply connected co-H spaces. Then f is a sum of iterated Whitehead products.*

Proof. Suppose $\dim X = k$ and $f: X \rightarrow G \vee H$ is given. Decompose $\Omega(G \vee H)$ as in theorem 3.6 and note that any product P_α of length k is at least k connected. Consequently the restriction of Ωf :

$$(\Omega X)^{k-1} \xrightarrow{\Omega f} \Omega(G \vee H)$$

factors through the product $\Omega \left(\prod_{\ell_\alpha \leq k} P_\alpha(k) \right)$ and the adjoint:

$$S \left[(\Omega X)^{k-1} \right] \rightarrow G \vee H$$

is a sum of iterated Whitehead products. However f is the composition of this map with the co-H space structure map:

$$X \rightarrow S \left[(\Omega X)^{k-1} \right]$$

which is a co-H map, so f is such a sum as well.

Proposition 3.8. *If G and H are simply connected, then there is a homotopy equivalence*

$$\phi: (G \vee H) \cup_W C(G \circ H) \rightarrow G \times H.$$

Proof. Since the composition

$$G \circ H \rightarrow \Omega G * \Omega H \rightarrow G \vee H \rightarrow G \times H$$

is null homotopic, there is an extension

$$C = (G \vee H) \cup C(G \circ H) \xrightarrow{\phi} G \times H.$$

The problem is to show that this map is a homotopy equivalence. We begin by observing that we can construct a right inverse ζ to $\Omega\phi$ as the sum of the loops on the inclusions of G and H into C :

$$\Omega G \times \Omega H \xrightarrow{\zeta} \Omega C \xrightarrow{\Omega\phi} \Omega G \times \Omega H$$

so $(\Omega\phi)_*: H_*(\Omega C) \rightarrow H_*(\Omega G \times \Omega H)$ is an epimorphism. We will complete the proof by showing that the rank of $H_k(\Omega C)$ is less than or equal to the rank of $H_k(\Omega G \times \Omega H)$. We will need several lemmas.

Lemma 3.9. *Write $\Omega G * \Omega H \simeq G \circ H \vee SQ$. Then the restriction:*

$$SQ \rightarrow \Omega G * \Omega H \xrightarrow{\omega} G \vee H \rightarrow C$$

is null homotopic

Proof. We first look at the homotopy commutative diagram

$$\begin{array}{ccccc} \Omega G * \Omega H & \xrightarrow{\omega} & G \vee H & \longrightarrow & G \times H \\ \downarrow & & \parallel & & \downarrow \pi_2 \\ G \rtimes \Omega H & \longrightarrow & G \vee H & \longrightarrow & H. \end{array}$$

Applying theorem 3a we see that the composition

$$SQ \rightarrow \Omega G * \Omega H \rightarrow G \rtimes \Omega H \rightarrow G \vee H$$

is a sum of maps γ_i factoring through $ad^i(H)(G) \rightarrow G \vee H$ for $i \leq 2$. By induction on i we see that

$$ad^i(H)(G) \rightarrow G \vee H \rightarrow C$$

is null homotopic for $i \geq 1$. This follows since ad^i factors

$$ad^i(H)(G) \rightarrow ad^{i-1}(H)(G) \vee H \xrightarrow{ad^{i-1} \vee 1} G \vee H \vee H \rightarrow G \vee H \rightarrow C.$$

It follows from 3.9 that the mapping cone of ω is homotopy equivalent to $C \vee S^2Q$. Recall that Ganea proved [Gan70] that given a fibration sequence $F \rightarrow E \rightarrow B$. One can construct a fibration sequence

$$F * \Omega B \rightarrow E \cup CF \xrightarrow{\pi} B$$

where π pinches the cone on F to a point. Apply this to the fibration sequence:

$$\Omega G * \Omega H \xrightarrow{\omega} G \vee H \rightarrow G \times H,$$

to obtain:

$$\Omega G * \Omega H * \Omega(G * H) \rightarrow C \vee S^2 Q \xrightarrow{\pi} G \times H.$$

It is possible that the map $\pi|_{S^2 Q}$ is nontrivial. However $\pi|_{S^2 Q}$ is the sum of the projections onto G and H , so it factors through C up to homotopy. Using such a factorization we can construct a homotopy equivalence:

$$\Gamma: C \vee S^2 Q \rightarrow C \vee S^2 Q$$

such that $\pi\Gamma|_{S^2 Q}$ is null homotopic. Replacing π with $\pi\Gamma$ does not alter the homotopy type of the fiber of π , so we can form the following diagram of fibrations:

$$\begin{array}{ccccc} S^2 Q \rtimes \Omega C & \longrightarrow & S^2 Q \rtimes \Omega C & & \\ \downarrow & & \downarrow & & \\ \Omega G * \Omega H * \Omega(G * H) & \longrightarrow & C \vee S^2 Q & \longrightarrow & G \times H \\ \downarrow & & \downarrow \pi_1 & & \parallel \\ K & \longrightarrow & C & \xrightarrow{\phi} & G \times H. \end{array}$$

The lefthand vertex fibration has a cross section since π_1 does, hence K is a co-H space and we have a splitting:

$$\Omega(\Omega G * \Omega H * \Omega(G \times H)) \simeq \Omega(S^2 Q \rtimes \Omega C) \times \Omega K.$$

Each space here is the loops on a co-H space, so the homologies are all tensor algebras. It follows that for each $i \geq 0$

$$\text{rank } H_i(\Omega G * \Omega H * \Omega(G \times H)) \geq \text{rank } H_i(S^2 Q \rtimes \Omega C).$$

We now calculate the Poincaré series for each of these spaces. Suppose $\mathcal{X}(G) = 1 + gt$ and $\mathcal{X}(H) = 1 + ht$ where g and h are polynomial in t with positive integral coefficient. We then have the following consequences:

$$\begin{aligned} \mathcal{X}(\Omega G) &= 1 + \frac{g}{1-g} \\ \mathcal{X}(\Omega H) &= 1 + \frac{h}{1-h} \\ \mathcal{X}(\Omega G * \Omega H) &= 1 + \frac{ght}{(1-g)(1-h)} \\ \mathcal{X}(G \circ H) &= 1 + ght \\ \mathcal{X}(SQ) &= 1 + ght \frac{g+h-gh}{(1-g)(1-h)} \\ \mathcal{X}(S^2 Q \rtimes \Omega C) &= 1 + ght^2 \frac{gth-gh}{(1-g)(1-h)} \cdot \mathcal{X}(\Omega C). \end{aligned}$$

On the other hand:

$$\mathcal{X}(\Omega G \times \Omega H) = 1 + \frac{g + h - gh}{(1 - g)(1 - h)}$$

Consequently

$$\mathcal{X}(\Omega G * \Omega H * (\Omega G \times \Omega H)) = 1 + \frac{ght^2(g + h - gh)}{(1 - g)^2(1 - h)^2}.$$

It follows that $\mathcal{X}(\Omega C) \leq \frac{1}{(1 - g)(1 - h)} = \mathcal{X}(\Omega G \times \Omega H)$ so $\text{rank } H_i(\Omega C) \leq \text{rank } H_i(\Omega G \times \Omega H)$ for all i and thus $(\Omega \phi)_\alpha$ is an isomorphism and hence ϕ is as well.

REFERENCES

- [AG95] D. Anick and B. Gray, *Small H spaces related to Moore spaces*, Topology **34** (1995), no. 4, 859–881.
- [Ani93] D. Anick, *Differential algebras in topology*, Research Notes in Mathematics, vol. 3, A K Peters, Ltd., Wellesley, MA, 1993, xxxvi+274 pp.
- [Ark62] M. Arkowicz, *The generalized Whitehead product*, Pacific J. Math **12** (1962), no. 1, 7–23.
- [CMN79] F. R. Cohen, J. C. Moore, and J. A. Neisendorfer, *Torsion in homotopy groups*, Ann. of Math. (2) **109** (1979), no. 1, 121–168.
- [Coh57] D. E. Cohen, *Products and carrier theory*, Proc. London Math. Soc. **7** (1957), no. 3, 219–248.
- [DT58] A. Dold and R. Thom, *Quasifassungen und unendliche symmetrische Produkte*, Ann. Math. **67** (1958), 239–281.
- [Gan70] T. Ganea, *Cogroups and suspensions*, Invent. Math. **9** (1970), 185–197.
- [Gra71] B. Gray, *A note on the Hilton–Milnor Theorem*, Topology **10** (1971), 199–201.
- [Gra88] ———, *On the iterated suspension*, Topology **27** (1988), no. 3, 301–310.
- [GT10] B. Gray and S. Theriault, *An elementary construction of Anick’s fibration*, Geometry and Topology **14** (2010), 243–276.
- [Hil59] P. J. Hilton, *Homotopy theory and duality*, mimeographed notes, 1959, Cornell University.
- [Nei09] J. Neisendorfer, *Algebraic methods in unstable homotopy theory*, New Mathematical Monographs, no. 12, Cambridge University Press, 2009.
- [The03] S. D. Theriault, *Homotopy decompositions involving the loops on a co-associative co-H space*, Canada J. Math. **55** (2003), no. 1, 181–203.
- [Whi41] J. H. C. Whitehead, *On adding relations to homotopy groups*, Annals of Math. **42** (1941), no. 2, 409–428.

DEPARTMENT OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE, UNIVERSITY OF ILLINOIS AT CHICAGO, 851 S. MORGAN STREET, CHICAGO, IL, 60607-7045, USA
E-mail address: brayton@uic.edu